

# Testosterone Effects on the Breast: Implications for Testosterone Therapy for Women

WORALUK SOMBOONPORN AND SUSAN R. DAVIS

*The National Health and Medical Research Council Centre of Clinical Research Excellence for the Study of Women's Health at the Jean Hailes Foundation and the Department of Obstetrics and Gynaecology, Monash University, Clayton, Victoria 3168, Australia*

Androgens have important physiological effects in women. Postmenopausal androgen replacement, most commonly as testosterone therapy, is becoming increasingly widespread. This is despite the lack of clear guidelines regarding the diagnosis of androgen insufficiency, optimal therapeutic doses, and long-term safety data. With respect to the breast specifically, there is the potential for exogenous testosterone to exert either androgenic or indirect estrogenic actions, with the latter potentially increasing breast cancer risk. In experimental studies, androgens exhibit growth-inhibitory and apoptotic effects in some, but not all, breast cancer cell lines. Differing effects between cell lines appear to be due primarily to variations in concentrations of specific coregulatory proteins at the receptor level. In rodent breast cancer models, androgen action is antiproliferative and proapoptotic, and is mediated via the androgen recep-

tor, despite the potential for testosterone and dehydroepiandrosterone to be aromatized to estrogen. The results from studies in rhesus monkeys suggest that testosterone may serve as a natural endogenous protector of the breast and limit mitogenic and cancer-promoting effects of estrogen on mammary epithelium. Epidemiological studies have significant methodological limitations and provide inconclusive results. The strongest data for exogenous testosterone therapy comes from primate studies. Based on such simulations, inclusion of testosterone in postmenopausal estrogen-progestin regimens has the potential to ameliorate the stimulating effects of combined estrogen-progestin on the breast. Research addressing this is warranted; however, the number of women that would be required for an adequately powered randomized controlled trial renders such a study unlikely. (*Endocrine Reviews* 25: 374–388, 2004)

- I. Introduction
- II. Testosterone Production and Metabolism
  - A. Biosynthesis of testosterone
  - B. Factors affecting circulating testosterone levels
- III. Mammary Epithelial Cell Proliferation and Apoptosis
- IV. Preclinical Studies of the Effects of Androgens in Breast Tissue
  - A. Breast cancer cell line studies
  - B. Animal studies
- V. Studies in Humans of the Effects of Androgens on Breast Cancer
  - A. Endogenous circulating testosterone and breast cancer
  - B. Exogenous testosterone therapy and breast cancer risk
  - C. Implications of the detection of the AR in human breast cancer
- VI. Aromatization and Breast Cancer Development
- VII. Should Androgens Be Included in Postmenopausal Hormone Therapy Regimens?
- VIII. Conclusion

Abbreviations: A, Androstenedione; AR, androgen receptor; CI, confidence interval; DHEA, dehydroepiandrosterone; DHEA-S, DHEA sulfate; DHT, dihydrotestosterone; DMBA, dimethylbenz(a)anthracene; ER, estrogen receptor; OR, odds ratio; PCOS, polycystic ovarian syndrome; PR, progesterone receptor; PSA, prostate-specific antigen; RR, relative risk.

*Endocrine Reviews* is published bimonthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

## I. Introduction

ANDROGENS HAVE IMPORTANT physiological effects in women. Not only are they the precursor hormones for estrogen biosynthesis in the ovaries and extragonadal tissues (1), but androgens act directly via androgen receptors (ARs) throughout the body. Androgen levels decline with increasing age in women before menopause (2, 3), and it is now accepted that many postmenopausal women experience a variety of physical symptoms secondary to androgen depletion, as well as physiological changes that affect their quality of life (4). Postmenopausal androgen replacement, most commonly as testosterone therapy, is becoming increasingly widespread. This is despite the lack of clear guidelines regarding the diagnosis of androgen insufficiency, optimal therapeutic doses, and long-term safety data. With respect to the breast specifically, there is the potential for exogenous testosterone to exert either androgenic or indirect estrogenic actions, with the latter potentially increasing breast cancer risk.

There is justifiable concern that combined oral estrogen plus progestin therapy significantly increases the risk of breast cancer in postmenopausal women (5–11). Although the underlying mechanism by which the development of breast cancer is increased in women taking combined hormone therapy is not understood, there is a considerable amount of evidence that androgens protect against estrogen's mitogenic and cancer-promoting effects on breast tissue. Labrie *et al.* (3) have recently reviewed the role of adrenal androgens in breast cancer growth with specific attention to dehydroepiandrosterone (DHEA). With the increasing in-

clusion of testosterone in hormonal regimens, the modulating effects of this steroid on the development of breast cancer therefore require consideration.

Thus, we have reviewed the published literature specifically pertaining to clinical studies of endogenous testosterone and testosterone therapy and breast cancer risk in premenopausal and postmenopausal women and examined the potential benefit or risk with regard to breast cancer of the administration of testosterone as part of hormone therapy.

## II. Testosterone Production and Metabolism

### A. Biosynthesis of testosterone

The term “androgens” refers to a group of 19-carbon steroid hormones that are associated with maleness and the induction of male secondary sexual characteristics. In women, androgens circulate in the concentration range nanomolar to micromolar, in contrast to the estrogens, the circulating concentrations of which are in the picomolar range. The major androgens in women include testosterone and dihydrotestosterone (DHT) because both have high binding affinity to the AR. Biosynthesis of the androgens takes place both in the adrenal and in the ovary and is modulated by two cytochrome P450 enzymes, P450 Scc, which catalyzes cholesterol side-chain cleavage, and P450 C<sub>17</sub>, which catalyzes 17-hydroxylation and 17–20 bond cleavage (17/20 lyase), which is required for the production of DHEA and androstenedione (A) from pregnenolone and progesterone, respectively. The further metabolism of androgens involves other important enzymes including 3 $\beta$ -hydroxysteroid dehydrogenase, catalyzing the conversion of pregnenolone to progesterone and DHEA to A, and 17 $\beta$ -hydroxysteroid dehydrogenase, which catalyzes the conversion of A to testosterone. DHEA secretion is acutely stimulated by ACTH (12, 13); however, DHEA sulfate (DHEA-S), which has a long plasma half-life, does not increase acutely after ACTH administration (14).

DHEA-S, DHEA, and A are considered to be proandrogens because they require conversion to testosterone to exhibit androgenic effects. Up to 25% of testosterone may be derived from the adrenal glands, 25% is derived from the ovary, and the remaining 50% is derived from peripheral conversion of the proandrogens, with A being the main precursor (15) (Fig. 1A). Circulating testosterone can be converted to DHT by 5 $\alpha$ -reductase, type 1 and type 2, or to estradiol by the aromatase enzyme. These conversions occur primarily in target tissues. DHT is a nonaromatizable androgen (16, 17) (Fig. 1B). Thus, androgens may exert biological effects by acting directly via the AR or indirectly after conversion to estrogen (17, 18).

### B. Factors affecting circulating testosterone levels

Under normal physiological conditions, only 1–2% of total testosterone circulates free, unbound to plasma protein. The rest is bound by SHBG and albumin, with SHBG binding 66% of total circulating testosterone (19). The binding affinity for steroids bound by SHBG is DHT > testosterone > androstenediol > estradiol > estrone (20). SHBG also weakly binds




FIG. 1. Androgen production and metabolism. A, The adrenal glands produce all proandrogens, whereas the ovaries produce only DHEA, A, and testosterone. In postmenopausal women, only about 25% of circulating testosterone is directly secreted by the ovaries. The rest, 50–75%, is formed largely from circulating precursors. B, Testosterone can be converted to DHT by 5 $\alpha$ -reductase type 1 and type 2 or to estradiol by the aromatase enzyme.

DHEA, but not DHEA-S (20). SHBG is a pivotal determinant of the bioavailability of sex steroids, and variations in the plasma levels of SHBG impact significantly on the amount of free and other bound sex steroids (20). Elevations in estradiol (as occurs during pregnancy, hyperthyroidism, and liver disease) cause a marked increase in SHBG levels, whereas hypothyroidism, obesity, and hyperinsulinemia are associated with decreased SHBG levels (21). Standard-dose oral estrogen, as used in hormone therapy, will increase SHBG with little or no effect seen with standard estradiol patch therapy (22).

In premenopausal women with regular menstrual cycles, there is a rise in testosterone and A in the late follicular phase of the menstrual cycle and in the luteal phase (23, 24). There is also a diurnal variation in testosterone in women with the peak in the morning (25). Zumoff *et al.* (2) showed lower mean 24-h values for total and calculated free testosterone among older *vs.* younger reproductive aged women ( $n = 33$  women). Most recently, a study of 149 healthy premenopausal women with regular cycles, no exogenous hormone therapy, and no complaint of low libido showed a statistically significant decline with age for free testosterone, DHEA-S, A, and DHT, each measured after organic solvent extraction by validated methodology (26). In the late reproductive years there is failure of the midcycle rise in free

testosterone that characterizes the menstrual cycle in young ovulating women (27). This occurs despite preservation of normal free testosterone levels at other phases of the cycle. The mean plasma concentrations of testosterone in women transiting the menopause are also significantly lower than those of younger ovulating women sampled in the early follicular phase (28). Across the perimenopausal period, neither A, DHT, or the ratio of total testosterone to SHBG (the free androgen index) appear to change acutely (19, 28). Controversy remains as to whether the postmenopausal ovary is a significant source of androgen production. Concentrations of testosterone in the ovarian vein of postmenopausal women have been shown to be higher than those in systemic venous blood, suggesting that the postmenopausal ovary continues to be an androgen-secreting organ (24). In addition, testosterone levels decrease in postmenopausal women after oophorectomy (29). However, Couzinet *et al.* (30) have proposed that the postmenopausal ovary is not a major source of androgens. Some postmenopausal women have elevated ovarian androgen production, *i.e.*, hyperthecosis, a well-established but poorly studied entity. It may well be that the androgen production of the postmenopausal ovary is variable. This variability requires further study along with associated factors.

### III. Mammary Epithelial Cell Proliferation and Apoptosis

Steroids and their nuclear receptors play crucial roles in the development and maintenance of normal functions of the human mammary gland. In addition to estrogen receptor- $\alpha$  (ER $\alpha$ ), estrogen receptor- $\beta$  (ER $\beta$ ), and progesterone receptors (PRs), ARs are present in both normal mammary tissue and breast cancer cell lines (31, 32). Hormone stimulation of mammary epithelial proliferation and apoptosis is important in tissue homeostasis. Deregulation of apoptosis can promote tumorigenesis as well as proliferation (33).

In premenopausal women, proliferation and apoptosis of normal breast epithelial tissue are higher in the luteal phase

of the menstrual cycle than in the follicular phase (34, 35). In the luteal phase, both estrogen and progesterone levels are maximal (36–38). Free testosterone and A levels peak during the middle-to-late follicular phase of the menstrual cycle and remain moderately elevated up through the midluteal phase (39).

*In vitro* studies of normal breast tissue are important to understand physiological regulation of the mammary gland by sex steroids. However, this has been hampered by experimental difficulties (40). One problem has been that normal human breast cells lose their steroid receptors and, hence, their hormone responsiveness almost as soon as they are isolated and placed into culture (40). Only one study of hormone-responsive primary cultures of breast epithelial cells reported that estradiol, but not progesterone, stimulated proliferation (41).

*In vitro* studies have consistently demonstrated that estradiol is a major mitogen in breast cancer cell lines (42–44). In breast cancer cell lines, cell death has been reported to be induced by estrogen deprivation (45) or antiestrogen (46). In contrast, *in vivo* evidence that supports a role for progesterone in cell proliferation in the breast has been difficult to reproduce *in vitro* (47).

### IV. Preclinical Studies of the Effects of Androgens in Breast Tissue

#### A. Breast cancer cell line studies

There is no *in vitro* evidence pertaining to the effects of androgens on normal human breast cells. Studies of the effects of androgens in various breast cancer cell lines predominantly support apoptotic and antiproliferative effects of androgens on the mitogenic effects of estrogens. However, divergent findings have been reported with differences according to the specific cell line, the androgen used, and its dose and estrogen status. The effects of testosterone and DHT in breast cancer cell lines are summarized in Table 1.

1. *Antiproliferative effects.* The antiproliferative effects depend on the following factors.

TABLE 1. Effects of testosterone and DHT on breast cancer cell lines

Study	Model	Type of androgen	Dose of androgen (M)	Main outcome measurement	Result
Boccuzzi <i>et al.</i> , 1994 (51)	MCF-7	DHT	$20^{-11}$ – $20^{-7}$	Proliferation	Biphasic effect: stimulation at a very high concentration; inhibition at concentration up to $20^{-8}$ M
Birrell <i>et al.</i> , 1995 (52)	MCF-7, ZR75-1, TD47-D, MDA-MB-453	DHT	$10^{-10}$ – $10^{-8}$	Proliferation	Stimulation in the MCF-7 and MDA-MB-453; no effect in MDA-MB-231 or BT-20, inhibition in T47-D and ZR-75-1
Ando <i>et al.</i> , 2002 (48)	MCF-7	Testosterone	$10^{-9}$ – $10^{-6}$	Proliferation	Inhibition
Ortmann <i>et al.</i> , 2002 (54)	MCF-7, T47-D, BT-20, MDA-MB 435S	DHT	$10^{-9}$ – $10^{-6}$	Proliferation	Inhibition
		Testosterone	$10^{-9}$ – $10^{-7}$	Proliferation	Inhibition in all four cell lines
Kandouz <i>et al.</i> , 1999 (58)	MCF-7, ZR75-1, T47-D	DHT	$10^{-9}$ – $10^{-7}$	Proliferation	Inhibition in all four cell lines
		DHT	$10^{-8}$	Apoptosis	Proapoptotic effect
Lapointe <i>et al.</i> , 1999 (61)	ZR-75-1	DHT	$10^{-9}$	Bcl-2 protooncogene	Down-regulation

*a. Estrogen status and type of androgen used.* Ando *et al.* (48) simulated the hormonal environment in pre- and postmenopausal women with an *in vitro* model utilizing the ER-positive breast cancer cell line MCF-7 exposed to DHEA, DHEA-S, androstenediol, testosterone, and DHT with or without estradiol. They found that DHEA-S and androstenediol stimulated the growth of MCF-7 cells in the absence of estradiol, but reduced cell proliferation in the presence of estradiol at 1 nmol/liter. This is consistent with the possibility of DHEA-S being converted to estrone and hence to estradiol, and androstenediol acting as a weak estrogen (49). Testosterone and DHT, at 1–100 nmol/liter, inhibited MCF-7 cell proliferation independently of the presence of estradiol. DHT alone, at 100 nmol/liter, consistently inhibited MCF-7 cell proliferation by 50% of the basal growth rate and counteracted estradiol-proliferative action by 68%. Normal circulating levels in women are approximately 0.5–2.3 nmol/liter for testosterone, and 0.2–0.8 nmol/liter for DHT. Thus, most of the concentrations for these hormones used in this and other studies are supraphysiological. Cell cycle analysis showed an enhanced number of cells in G<sub>0</sub>/G<sub>1</sub> phase after 6 d of DHT treatment. Moreover, upon prolonged DHT exposure, a markedly increased AR content, mostly translocated into the nucleus, was observed. The inhibitory effect of DHT on cell proliferation was lost when the cells were treated with the AR antagonist, hydroxyflutamide (48). In cotransfection experiments, overexpression of the AR decreased estradiol-induced signaling (48). This was amplified by treatment with DHT but lost with the addition of hydroxyflutamide. When the cells were cotransfected with a mutant AR, inhibition of estradiol-induced signaling did not occur. Thus, direct androgen action appears to antagonize MCF-7 proliferation induced by estradiol, and this seems to be related to the inhibitory effects of the AR on estradiol genomic action (48). These experimental results are consistent with those of Birrell *et al.* (50) who proposed that the therapeutic action of medroxyprogesterone acetate in breast cancer may be partially mediated by the AR.

*b. Androgen concentration.* The effects of DHT appear to be concentration dependent. Boccuzzi *et al.* (51) reported that, at an extremely high concentration (200 nmol/liter), DHT stimulated MCF-7 cell growth through an ER-mediated mechanism, whereas lower concentrations of DHT were inhibitory.

*c. Type of breast cancer cell line.* The effects on proliferation *in vitro* vary according to the androgen administered and the breast cancer cell line studied (52, 53). At concentrations of 1 nmol/liter for 10 d, which is close to the normal female physiological range, Birrell *et al.* (52) reported varying stimulatory and inhibitory effects of DHT on differing breast cancer cell lines that simply could not be explained by varying hormone receptor status. The two cell lines that had no hormone receptors did not respond to treatment. DHT stimulated both the ER-positive MCF-7 cells and the ER-negative MDA-MB-453 cells. Treatment with 100-fold excess of hydroxyflutamide reversed the effects of DHT in each of the cell lines. In the same study (data not shown) the synthetic non-metabolizable androgen mibolerone had effects similar to those of DHT, with the exception that hydroxyflutamide only

partially reversed the growth-stimulatory effects of this treatment on MCF-7 and MDA-MB-453 cells. Hydroxyflutamide only partially reversed the inhibitory effects of DHT on ZR-75-1 cultures, whereas AR antisense oligonucleotides reversed the growth-inhibitory action of mibolerone in this cell line.

These observations support the theory that AR expression is a necessary requirement for androgenic effects on breast cancer cell proliferation, but that the absolute levels of AR (as well as ER and PR) in cell lines are associated with neither a specific stimulatory nor inhibitory proliferative responses (52). It is likely, therefore, that additional cellular factors or the structure of the AR determine whether breast cancer cell proliferation is stimulated or inhibited in the presence of androgen (53).

In contrast to the above, Ortmann *et al.* (54) reported dose-dependent inhibition with androgens of four cell lines that each stained positively for the AR. Included among these was the BT-20 cell line, which was reported by two other groups to be AR negative (52, 55). According to proliferation kinetics, they observed a significant ( $P < 0.05$ ) dose-dependent inhibition of cell growth by both testosterone and DHT. In the ER-negative cell lines, BT-20 and MDA-MB 4<sup>35</sup>S testosterone was a more potent inhibitor of cell proliferation than DHT ( $P < 0.05$ ), whereas in the ER-positive cells lines, MCF-7 and T47-D, stronger inhibition of proliferation was achieved with DHT than with testosterone. They proposed that partial transformation of testosterone to estrogen in ER-positive cells might be an explanation for this effect (54).

Prostate-specific antigen (PSA) is a new favorable prognostic indicator for women with breast cancer (56). Immunoreactive PSA has been reported in 30% of breast cancers and has been associated with both earlier stage disease and longer relapse-free survival (56). KLK3 (which encodes PSA) and KLK2 (encoding human kallikrein 2) are both known to be androgen regulated, but respond differentially to androgens when studied in different human breast cancer cell lines (57). Research into the various factors affecting the production of these two proteins, according to the breast cancer cell line studied, gives insight into how androgen treatment may affect different cell lines differently.

Initial experiments demonstrated that the differential androgen induction of PSA and human kallikrein 2 was not directly related to the level of AR expression or to mutations within the coding region (55). Because the action of a steroid receptor at a given promoter may be modulated by various coregulatory proteins (coactivators/corepressors), Magklara *et al.* (55) examined the expression of various known coregulatory proteins in the different breast cancer cell lines (BT-474, T-47D, ZR75-1, MCF-7, MFM-223 and BT-20). They found that the mRNA levels of steroid receptor coactivator 1, a known coactivator of the activation function 1 domain of the AR, were highest in the breast cancer cell lines with the greatest PSA production and lowest in the cell lines that secreted less PSA. This raises the possibility that the relative levels of specific coactivators/corepressors might differentially modulate AR transcriptional activity within the promoter/enhancer region of KLK2 and KLK3 of different breast cancer cell lines (55).

Thus, the ultimate effects of testosterone and DHT at the

tissue level may not be influenced simply by absolute ligand concentrations, but by the relative concentrations of specific coregulatory proteins unique to each cell line. Hence, the effects of DHT differ between different breast cancer cell lines, despite the consistent presence of ARs. In the AR-positive human breast cancer cell lines, T47-D and ZR-75-1, DHT has been reported to be proapoptotic, with maximal effects at 10 nmol/liter (58). These findings are in line with the growth-inhibitory effects reported by Birrell *et al.* (50) in these two cell lines.

**2. Apoptotic effects.** The effects of androgens on the expression of two genes known to influence apoptosis, Bcl-2 and Bax, have been investigated. Bcl-2 is able to counteract apoptosis induced by numerous stimuli such as UV light,  $\gamma$ -radiation, heat shock, and chemotherapy and thus is considered antiapoptotic (59), and Bax is a proapoptotic gene (60). Lapointe *et al.* (61) reported that DHT down-regulated Bcl-2 protooncogene levels via an AR-mediated mechanism in the ZR75-1 breast cancer cell line in either the presence or absence of  $17\beta$ -estradiol. This is consistent with the inhibitory effect of DHT in this cell line. Inhibition by DHT was completely prevented by coincubation with the pure antiandrogen hydroxyflutamide (61). Xie *et al.* (60) studied Bcl-2 and Bax expression in the Noble rat model, which they had established to explore the mechanisms of hormonal mammary carcinogenesis. They observed that Bcl-2 was strongly expressed in most of the mammary tumor cells, and that when animals were treated with  $17\beta$ -estradiol, the mammary epithelial cells expressed higher levels of Bcl-2. Bax immunoreactivity was weak in mammary tumor cells but strongly expressed in adjacent normal or hyperplastic ductal structures. Treatment with testosterone, either alone or in combination with estrogen, gave rise to high levels of Bax expression in “pre-malignant” mammary glands. This supports the hypothesis that testosterone may oppose the mitogenic action of estrogen in the breast by promoting apoptosis (62).

Taken together, androgens exhibit growth-inhibitory and apoptotic effects in some, but not all, breast cancer cell lines. Differing effects between cell lines appear to be primarily due to variations in concentrations of specific coregulatory proteins at the receptor level. In rodent breast cancer models, androgen action is antiproliferative and proapoptotic, and mediated via the AR, despite the potential for testosterone and DHEA to be aromatized to estrogen.

## B. Animal studies

**1. DHEA.** The effects of DHEA have been renewed in detail recently (3). In summary, data from *in vivo* studies support a primarily inhibitory effect of testosterone on the proliferative effects of estrogen. Dimethylbenz(a)anthracene (DMBA)-induced mammary carcinoma in the rat is a commonly used model for *in vivo* studies of the prevention and treatment of breast cancer. Labrie and associates (63–65) and Lubet *et al.* (66) have used this model extensively to study the effects of DHEA and have consistently reported an inhibitory effect of DHEA on mammary carcinoma de-

velopment. Treatment with SILASTIC (Dow Corning Corp., Midland, MI) implants of DHEA leading to serum DHEA levels comparable to those observed in normal adult women ( $7.1 \pm 0.6$  nmol/liter and  $17.5 \pm 1.1$  nmol/liter) progressively inhibited the development of DMBA-induced mammary carcinoma in the rat (63). Luo *et al.* (64) also demonstrated inhibition in DMBA-induced mammary carcinoma development: 279 d after DMBA administration, the incidence of mammary carcinoma had decreased from 95% in control animals to 73% ( $P < 0.05$ ), 57% ( $P < 0.01$ ), and 38% ( $P < 0.01$ ) at the daily percutaneous doses of 5, 10, and 20 mg of DHEA, respectively. Moreover, the mean tumor number and the mean tumor area per tumor-bearing animal were also reduced by the same treatments. Similar outcomes have been reported in a *N*-methyl-*N*-nitrosourea-induced rat mammary tumor model with DHEA therapy (65), and a suppressive effect of DHEA has been demonstrated on the growth of estrone-stimulated sc tumor xenografts formed by AR-positive ZR-75-1 cells in ovariectomized nude mice (66).

**2. Testosterone and DHT.** Table 2 summarizes the effects of testosterone and DHT in animal studies. There is both direct and indirect evidence of the inhibitory effects of testosterone and DHT on DMBA-induced mammary carcinoma in ovariectomized rats (67, 68). DHT therapy reduced the number of progressing tumors from 69.2 to 29.2% in estradiol-treated animals, and the number of tumors that completely regressed increased from 11.5 to 33.3% in the same groups of animals (67). Simultaneous treatment with the antiandrogen flutamide completely prevented the inhibitory effect of DHT on tumor growth (67). DHT exhibited similar tumor-suppressing effects on ZR-75-1 breast cancer cells implanted into ovariectomized athymic mice in either the presence or absence of exogenous estradiol (69).

Zhou *et al.* (31) explored the effects of estrogen, progesterone, and testosterone on normal mammary epithelial cell proliferation and steroid receptor gene expression in the ovariectomized primate mammal. They showed that estrogen therapy alone significantly increased mammary epithelial proliferation approximately 6-fold and significantly increased the mammary epithelial level of ER $\alpha$  mRNA. Progesterone administration did not modify the proliferative effects of estradiol significantly. When given concurrently, testosterone reduced estradiol-induced epithelial proliferation by approximately 40% and entirely abolished the estradiol-induced augmentation of ER $\alpha$  gene expression. However, testosterone levels achieved in this study were supraphysiological. Zhou *et al.* (31) also investigated the effects of tamoxifen and found that it caused a 3-fold increase in mammary epithelial proliferation, measured by Ki67, but a decrease in ER $\alpha$  gene expression below placebo level. AR mRNA levels detected by *in situ* hybridization were not altered by estradiol alone, but were significantly reduced by estradiol plus testosterone or tamoxifen treatment. Thus, testosterone induced down-regulation of mammary epithelial proliferation and ER $\alpha$  gene expression. This suggests that the addition of testosterone might reduce the risk of breast cancer associated with estrogen-progestin therapy in postmenopausal women. In addition to the parallel effects of tamoxifen

TABLE 2. Effects of testosterone and DHT in animal studies

Study	Model	Intervention	Main outcome measurement	Result
Dauvois <i>et al.</i> , 1989 (67)	DMBA-induced mammary carcinoma in rats	E vs. E + DHT	The number of progressing tumors The number of complete responses	Significant decrease with E + DHT vs. E alone Significant increase with E + DHT vs. E alone
Dauvois <i>et al.</i> , 1991 (69)	ZR-75-1 breast cancer cells implanted into ovariectomized athymic mice	E vs. E + DHT	The number of progressing tumors The number of complete responses	Significant decrease with E + DHT vs. E alone Significant decrease with E + DHT vs. E alone
Jayo <i>et al.</i> , 2000 (71)	Rat	Oral contraceptive vs. oral contraceptive + methyltestosterone	Mammary epithelial proliferation	Significant decrease with oral contraceptive + testosterone vs. oral contraceptive alone
Zhou <i>et al.</i> , 2000 (31)	Ovariectomized rhesus monkeys	E; E + progesterone; E + T	Mammary epithelial proliferation	Significant decrease with E + T vs. others
Dimitrakakis <i>et al.</i> , 2003 (70)	Normal-cycling rhesus monkeys Ovariectomized rhesus monkeys	Flutamide vs. none E; E + progesterone; E + T; vehicle	Mammary epithelial proliferation Mammary epithelial proliferation ER $\alpha$ /ER $\beta$ MYC expression	Significant increase in flutamide Significant decrease with E + T vs. E or E + progesterone Reversal of ER $\alpha$ /ER $\beta$ Significant reduction

E, Estradiol; T, testosterone.

on primate mammary epithelial sex steroid receptor gene expression, Zhou *et al.* (31) also demonstrated that tamoxifen, like testosterone, reduced apolipoprotein D mRNA levels and increased IGF binding protein 5 expression in the primate mammary gland. These findings support AR-mediated effects of tamoxifen.

Subsequently, Dimitrakakis *et al.* (70) have investigated the effects of ovarian steroids in physiological doses on the mammary epithelial proliferation in ovariectomized rhesus monkeys. They studied four groups treated with placebo, estradiol alone, estradiol plus progesterone, or estradiol plus testosterone. Circulating estradiol levels were similar in the three active treatment groups, and testosterone levels were physiological. The mammary epithelial proliferation index was measured using Ki67 immunoreactivity. Estradiol alone and estradiol plus progesterone resulted in a significantly increased mammary epithelial proliferation index compared with placebo controls by approximately 3.5-fold, whereas the estradiol plus testosterone combination did not increase the proliferation index above control values (70). In addition, they found a significant reduction in mammary epithelial ER $\alpha$  and increase in ER $\beta$  expression in estradiol plus testosterone groups compared with estradiol alone. This effect of testosterone resulted in a reversal of the ER $\alpha$ /ER $\beta$  ratio, which was approximately 2.5 in the estradiol-treated group and approximately 0.7 in the estradiol-testosterone group. Moreover, testosterone treatment was associated with an approximate 50% reduction in mammary epithelial MYC expression, an estrogen-responsive gene, compared with the estradiol- and estradiol-progesterone-treated groups ( $P = 0.05$ ), suggesting that the antiestrogenic effects of testosterone at the mammary gland involve alteration in ER signaling to MYC (70). They further investigated the importance of endogenous testosterone in intact, cycling monkeys by

studying treatment with either placebo or the AR antagonist flutamide. The mammary epithelial proliferation index increased 2-fold after treatment with flutamide alone ( $P = 0.03$ ). This suggests that testosterone may serve as a natural endogenous protector of the breast and may limit mitogenic and cancer-promoting effects of estrogen on the mammary epithelium (70). Jayo *et al.* (71) also reported that oral contraceptive therapy plus methyltestosterone in rats causes significant suppression of epithelial cell proliferation, a reduction in the number of proliferating cell nuclear antigen-labeled cells, and an increase in the number of PR-labeled cells. However, there have been no studies of the effects of testosterone plus estrogen-progestin therapy on breast epithelial cell proliferation in women.

In ovariectomized 12-month-old rats, DHT stimulated mammary gland lobulo-alveolar ductal growth (72). Medroxyprogesterone acetate induced the same effect, consistent with the findings of Birrell *et al.* (50) that androgens and medroxyprogesterone acetate share a similar mechanism of action in the breast. DHEA also stimulated lobulo-alveolar development, which was unaffected by the antiestrogen EM-800 but almost completely prevented by cotreatment with the antiandrogen flutamide (47, 72).

Taken together, available *in vivo* data indicate that in estrogen-replete normal breast tissue, androgens diminish estrogen-induced breast epithelial proliferation and abolish estrogen-induced ER $\alpha$  gene expression in the primate. In the absence of estrogen, androgen action mimics that of progestins in the rat mammary gland.

Thus results from the studies *in vitro* and *in vivo* suggest that testosterone may serve as a natural endogenous protector of the breast and limit mitogenic and cancer-promoting effects of estrogen on mammary epithelium. However,

these are surrogate endpoints and hence the need to consider what is known from studies in women.

## V. Studies in Humans of the Effects of Androgens on Breast Cancer

### A. Endogenous circulating testosterone and breast cancer

1. *Issues in clinical trials.* Findings from case-controlled studies of the relationship between endogenous testosterone levels and breast cancer risk do not necessarily translate to women treated with exogenous testosterone. The former address endogenous androgen production, which in some women may be pathophysiological. In contrast, postmenopausal testosterone therapy is administered to women who have low testosterone levels due to low production and who are usually also treated with exogenous estrogen. Furthermore, if an association is found between endogenous circulating testosterone and breast cancer, it does not necessarily signify a causal relationship. Total testosterone, although the most common measure for clinical studies to date, does not yield specifically meaningful information about actual tissue androgen exposure. It is widely accepted that free testosterone is the strongest indicator of tissue androgen exposure and that variations in SHBG levels in women can have dramatic effects on free testosterone levels (4, 73, 74). In addition, a single value may be inadequate to assess true tissue exposure because testosterone levels vary in response to diurnal rhythms (25). Stress is also an important confounder in cross-sectional studies, because stress itself affects testosterone levels (75–78). Whether even free testosterone is a meaningful indicator of tissue androgen exposure remains controversial. Labrie *et al.* (79, 80) proposed that the major proportion of androgenic effects in women are derived from an intracrine mode of action, which will not be detected by measurement of circulating testosterone or DHT. Regarding the type of study, a clear inference of effect cannot be drawn from cross-sectional data because such research cannot provide an appropriate time sequence of exposure and outcome.

Consideration also needs to be given not only to what has been measured but also the effects of storage and the sensitivity of the assay methodology used. For example, with long-term cryopreservation, testosterone has been shown to increase by 5% per year of blood storage at  $-20\text{ C}$  (81). This may not apply to storage at much colder temperatures (82).

No rapid, simple assay of total testosterone has been shown to produce reliable results in women with low to normal testosterone levels. Direct testosterone immunoassays are limited by “noise” from assay interference and by cross-reactivity with other steroids, which become worse at low testosterone concentrations (83). Inclusion of an organic solvent extraction step when measuring total testosterone will increase the assay specificity, and if combined with chromatographic separation of testosterone from interfering steroids, a reliable result can be obtained. The gold standard methodology for measurement of free testosterone is considered to be equilibrium dialysis. Measurement of free testosterone by analog assay is notoriously unreliable, particularly at the lower end of the normal female range and is not recommended for use (83).

Finally, because estrogen is considered a strong risk factor for breast cancer, to draw any conclusion about an association between testosterone and breast cancer, a statistical method to adjust for the estrogen effect must be employed.

2. *Testosterone levels and breast cancer in premenopausal women.* Two cross-sectional studies have investigated the relationship between total testosterone and breast cancer risk in premenopausal women and have yielded inconsistent results (Table 3). Secreto *et al.* (84) reported an age-adjusted relative risk (RR) for high *vs.* low levels of serum total testosterone of 3.4 [95% confidence interval (CI), 1.6–7.3] and for urinary testosterone of 2.1 (95% CI, 0.9–4.8) for cases ( $n = 63$ ) *vs.* controls ( $n = 70$ ). Study samples were collected between cycle d 18 and d 21 irrespective of cycle length or whether ovulation had or had not occurred. The association was observed only in women whose samples were collected 5–9 d before the next menses (a period corresponding to the mid-luteal phase) and 10 or more days before the next menses. There was no positive association for women whose blood and urine were collected within 4 d of the next menses and who thus had cycles lasting 25 d or less. The authors’ interpretation of these findings was that higher testosterone levels were detectable in cases only in the follicular or early luteal phases; however, these phases are the times of highest testosterone levels during the normal cycle (39). They also reported that high testosterone was characteristic of breast cancer patients with long menstrual cycles ( $>28$  d) and negligible in women with short ones ( $<28$  d) but that low SHBG levels appeared to be a protective factor for breast cancer. Thus, the high total testosterone measured corresponded to periovulatory hormone production but was not abnormally high across the cycle in premenopausal women with breast cancer, consistent with the fact that SHBG was not abnormally low in patients with breast cancer.

The most recent cross-sectional study involved 171 premenopausal women with breast cancer and 170 controls matched by age (85). No significant difference by odds ratio (OR) for breast cancer between high and low testosterone was found when data were adjusted for confounding factors. Overall, no conclusions can be made about testosterone and breast cancer in premenopausal women based on these two studies.

Two prospective case-control studies in which total testosterone was measured provide more consistent results (Table 3). Wysowski *et al.* (86) found no statistical associations between serum hormone levels, including total testosterone, in 17 women diagnosed with breast cancer 8–132 months after blood was drawn, each matched to four controls. Similarly, in a study of 62 premenopausal women with breast cancer and 182 controls, Thomas *et al.* (87) found no statistical difference for total testosterone between the groups.

3. *Polycystic ovarian syndrome (PCOS) and incidence of breast cancer.* In premenopausal women, PCOS is characterized by infertility, hyperandrogenism, and obesity. Concentrations of testosterone, A, and DHEA-S, and the calculated free androgen index (total testosterone in nanomoles per liter / SHBG in nanomoles per liter  $\times 100$ ) are significantly higher in women with PCOS regardless of hirsutism (88). This syn-

TABLE 3. Epidemiological studies of the association between plasma testosterone levels and risk of breast cancer in premenopausal women: study size, characteristics, and summary results

Study	Trial type	Study characteristics	Comparison made	Case/control (no. of women)	Type of T measurement	OR (95% CI)
Secreto <i>et al.</i> , 1989 (84)	Cross-sectional	Matched to age ( $\pm$ 6 months)	Top to bottom quartile	75/150	Total T	2.1 <sup>a</sup>
Yu <i>et al.</i> , 2003 (85)	Cross-sectional	Matched to age	Top to bottom tertile	171/170	Total T	3.4 (1.6–7.3) <sup>b</sup> 1.9 (1.0–3.7) <sup>c</sup> 2.01 (0.96–4.2) <sup>d</sup>
Wysowski <i>et al.</i> , 1987 (86)	Prospective	7-yr follow-up; matched to race, age, and time since last menstrual period	Mean values for cases <i>vs.</i> controls <sup>e</sup>	17/68	Total T	Not applicable
Thomas <i>et al.</i> , 1997 (87)	Prospective	Matched to age, year of blood collection, and no. of years postmenopausal	1U increase in the natural log of hormone concentration	61/179	Total T	1.2 (0.6–2.4) <sup>f</sup>

T, Testosterone.

<sup>a</sup> Age adjusted.<sup>b</sup> Adjusted for occupation and no. of children.<sup>c</sup> Unadjusted.<sup>d</sup> Adjusted for waist-hip ratio, age at first live birth, total caloric intake, fibroadenoma, and SHBG.<sup>e</sup> Mean comparison resulted in no statistically significant difference of testosterone levels between cases and controls.<sup>f</sup> Unadjusted OR.

TABLE 4. Epidemiological studies of the association between plasma testosterone levels and risk of breast cancer in women with PCOS: study size, characteristics, and summary results

Study	Trial type	Study characteristics	Comparison made	RR (95% CI)
Coulam <i>et al.</i> , 1983 (89)	Prospective cohort	Clinic-based study; 1270 subjects diagnosed as having chronic anovulatory syndrome were followed during 1935–1980	Observed incidence of breast cancer <i>vs.</i> expected incidence based on standard population	1.5 (0.8–2.6)
Gammon and Thompson, 1991 (91)	Case-control	Population-based study; 4730 women with breast cancer and 4688 control women aged 20–54 yr	OR	0.5 (0.3–0.9) <sup>a</sup>
Anderson <i>et al.</i> , 1997 (90)	Prospective cohort	Population-based study; 34,835 women at risk, age 55–69 yr, were followed during 1986–1992	Incidence of breast cancer among women with Stein-Leventhal syndrome <i>vs.</i> incidence among women without this disease in the same cohort	1.2 (0.7–2) <sup>b</sup> 1.0 (0.5–1.8) <sup>c</sup>

<sup>a</sup> Age-adjusted OR for age.<sup>b</sup> Adjusted for age.<sup>c</sup> Adjusted for age, age at menarche, age at first pregnancy, parity, oral contraceptive use, hormone replacement therapy, body mass index, waist-to-hip ratio, benign breast disease, and family history of breast carcinoma.

drome is a useful model of the effects of long-term exposure to hormone imbalance. An increased risk of endometrial cancer has been documented in women with this condition (89). Table 4 summarizes the studies of PCOS and breast cancer risk. Despite hyperandrogenism and long-term exposure to unopposed estrogen, the risk of breast cancer is not increased in women with PCOS (89, 90). In fact, Gammon and Thompson (91) have reported an age-adjusted OR for breast cancer in women with this syndrome of 0.52 (95% CI, 0.32–0.87). Although this risk reduction might be related to the hyperandrogenemia of this condition, a cause and effect cannot be established.

#### 4. Testosterone and breast cancer in postmenopausal women

*a. Cross-sectional studies.* In their cross-sectional studies, Lipworth *et al.* (92) measured hormone levels in cases 1 wk

post surgery, and Secreto *et al.* (93) collected blood and urine from women after diagnosis, but before surgery. Each group felt their protocol minimized the influence of stress, but this is questionable. Both studies reported only total testosterone, not free or bioavailable testosterone. Secreto's group (93) reported that breast cancer patients with elevated urinary testosterone levels at the time of diagnosis showed a dramatic decrease in the excretion levels of this hormone after bilateral ovariectomy, and that histological examination of the excised ovaries revealed hyperplasia of interstitial cells in all hyperandrogenic patients. No such change was observed in patients with normal testosterone levels at the time of diagnosis preovariectomy (94). This implies that the hyperandrogenic postmenopausal women who developed breast cancer in Secreto's study had ovarian pathophysiology. Lipworth *et al.* (92) reported no increase for breast cancer



with higher testosterone levels after adjustment for age, residence, and hormonal factors (OR, 0.48; 95% CI, 0.12–1.90), whereas Secreto *et al.* (93) reported a positive association after adjustment for occupation and number of children (OR, 2.7; 95% CI, 1.1–6.7). Secreto *et al.* (93) also measured DHT in postmenopausal women with breast cancer *vs.* controls and found no significant difference between the two groups.

*b. Prospective case-control studies.* Ten prospective case-control studies have been undertaken (81, 82, 86, 95–101). Study designs, characteristics, and results of these studies are listed in Table 5. All met the appropriate requirements for prevention of biochemical measurement bias, and case-control identification among these studies was similar. To address intraindividual variability of hormone levels with time, Berrino *et al.* (95), Hankinson *et al.* (102), and Thomas *et al.* (81) reported intraclass correlation coefficients between two blood samples. Several groups measured only total testosterone (81, 82, 86, 97, 98, 100). Overall, three of the studies did not demonstrate any significant associations without adjustment for estradiol (82, 86, 98). Dorgan *et al.* (96) reported the RR for breast cancer for women in the highest *vs.* the lowest quartile of total testosterone levels as being 6.2 (95% CI, 2.0–19.0). However, they did not control for estradiol in their calculations. Positive associations between breast cancer and total testosterone that were no longer significant after adjustment for estradiol were demonstrated by three groups (81, 97) or estrone (101). In contrast, Manjer *et al.* (100) reported an association between high total testosterone and breast cancer risk after adjustment for estradiol. Only two groups (95, 99) measured free testosterone. Cauley *et al.* (99) measured free testosterone by equilibrium dialysis, considered to be the most accurate methodology, in 97 cases and 244 controls and reported no association between free testosterone and breast cancer after adjustment for estradiol. In contrast Berrino *et al.* (95) measured free testosterone by RIA in 25 breast cancer patients and 100 controls and reported a significant association between higher free testosterone levels and breast cancer after adjustment for estradiol.

A reanalysis of prospective studies (103) reported that the RRs associated with a doubling of total testosterone levels were 1.37 (95% CI, 1.15–1.65) for the three studies incorporating a purification step in their testosterone assay (98, 99, 104) and 1.44 (95% CI, 1.21–1.72) for the four studies that used a direct testosterone assay (81, 95–97). However, a subgroup analysis to determine the association between breast cancer and free testosterone levels was not undertaken. Thus, evidence from clinical studies that the free fraction of testosterone is an independent risk factor for breast cancer is lacking.

### B. Exogenous testosterone therapy and breast cancer risk

Three observational studies have addressed the use of testosterone therapy and breast cancer risk (105–107). Unfortunately, the primary aim for two of these studies was not testosterone supplementation and breast cancer risk; consequently, they each had only a small sample size for this subgroup analysis. Brinton *et al.* (108) undertook a case-control study of postmenopausal estrogen use and breast

cancer risk. A subgroup analysis in this study of 25 patients and 29 controls showed no significant increase in risk with oral methyltestosterone in combination with conjugated equine estrogen (RR, 1.05; 95% CI, 0.6–1.8) (108). In contrast, Ewertz (109) studied the effects of the ever-use of im injections containing estradiol-testosterone (2.5 mg estradiol plus 50 mg testosterone or 5.0 mg estradiol plus 100 mg testosterone) given at a recommended interval of 3–7 wk in a subgroup analysis. This specific therapy was used for 56 of 1694 patients and 21 of 1705 controls. An RR of 2.3 (95% CI, 1.37–3.88) was reported (109). In the same study, there was no risk for triple combination of estrogen, progestin, and testosterone (RR 1.26; 95% CI, 0.58–2.74) (109). A recent retrospective analysis of 511 Australian women treated with conventional estrogen therapy plus 50–150 mg testosterone implants with a mean follow-up of  $5.7 \pm 2.5$  yr, but no control group, reported a breast cancer incidence of 240 per 100,000 women years (107). This was reported as equivalent to the incidence in the general population determined by the state cancer registry.

### C. Implications of the detection of the AR in human breast cancer

ARs are found in more than 50% of breast tumors (110), and the significance of ARs in breast cancer has been extensively explored. With regard to nodal metastasis, Soreide *et al.* (111) reported that when the median value of AR is taken as cut-off (50.5 pmol/g), a lower AR content is an independent predictor of the likelihood of axillary metastases ( $P = 0.001$ ). AR amount, however, did not reveal any significant prognostic information concerning relapse-free survival. There appears to be no significant association between AR expression and the degree of differentiation of ductal carcinoma *in situ* (112).

In addition to the amount of AR, correlations between the repeat length of the CAG sequence in the AR and total risk of breast cancer, age at diagnosis, recurrence after surgery, and aggressive growth have been reported. CAG repeat length is associated with a decreased ability to activate AR-responsive genes (113). More CAG repeats in the AR gene have been associated with an earlier onset of breast cancer among BRCA1 mutation carriers (114). However, Kadouri *et al.* (115) found no significant association between the number of CAG and GGC repeats in the AR and breast cancer risk in either BRCA1/2 carriers or the general population if attention was restricted to Ashkenazi Jewish carriers, or only to BRCA1 or BRCA2 carriers. One explanation for the discrepancy is sample size in that the larger number of study subjects with the BRCA1 mutation in the former study (165 with and 139 without breast cancer) could provide a statistical difference rather than smaller study subjects with BRCA1 and BRCA2 mutations (122 with and 66 without breast cancer). Therefore, an effect of the AR repeat length on BRCA1 penetrance cannot be excluded. However, this remains inconclusive for BRCA2.

The relationship between the number of CAG and GGC repeats has also been evaluated in a population-based study conducted in 524 patients and 461 controls for their relationships to breast cancer risk (116). This study suggested a

TABLE 5. Prospective studies of the association between plasma testosterone levels and risk of breast cancer in postmenopausal women

First author, year (Ref.)	Definition of study case	Mean or median time to diagnosis (yr)	No. of cases/controls	Mean or median age (yr)	Type of T assay	Ratio of control subjects to case patients	Unadjusted effect ratios (95% CI)	Adjusted RR by other variables	Effect ratios <sup>a</sup> (95% CI) <sup>b</sup>
Wyowski, 1987 (86)	Excluded new cases diagnosed within 6 months of recruitment	2.3	39/156	61.0 <sup>c</sup> 61.0 <sup>c</sup>	Total T	Matched 4:1	N/A	N/A	N/A <sup>d</sup>
Garland, 1992 (82)	All	9.0	31/287	65.3 66.6	Total T	Full cohort	<sup>e</sup>	Age	1 <sup>f</sup>
Berrino, 1996 (95)	All	3.5	67/264	59.4 54.9	Total T, free T	Matched 4:1	5.9 (1.6–21.9)	Age Estradiol	5.7 (1.5–22.2) 5.9 (1.2–29.3)
Dorgan, 1996 (96)	All	2.9	71/133	61.0 62.0	Total T	Matched 2:1	3.7 (1.4–10.0) <sup>g</sup>	Years since menopause, height, weight, parity, family history of breast cancer	6.2 (2.0–19.0)
Thomas, 1997 (81)	All	7.8	61/179	58.6 58.5	Total T	Matched 3:1	2.4 (1.0–5.7)	Age at menarche, parity, BMI, years postmenopausal, Estradiol	Not different 0.8 (0.3–2.4) 1.2 (0.4–3.5)
Zeleniuch-Jacquotte, 1997 (97)	Excluded new cases diagnosed within 6 months of recruitment	2.7	85/163	59.2 59.1	Total T	Matched 4:1	2.7 (1.1–6.8)	Total estradiol, SHBG-bound estradiol	
Hankinson, 1998 (98)	All	2.4	155/310	62 <sup>c</sup> 62 <sup>c</sup>	Total T	Matched 2:1	1.34 <sup>f</sup>	BMI at age 18, family history of breast cancer, age at menarche, parity/age at first birth, and past HT	1.4 (0.7–2.7)
Cauley, 1999 (99)	All	3.2	97/243	70.9 71.8	Total T, free T	Subcohort	2.7	Estradiol Age, BMI, age at menarche, first birth, menopause, family history of breast cancer, physical activity, surgical menopause, alcohol	1.1 (0.5–2.3) 3.3 (1.1–10.3)
Manjer, 2003 (100)	All	5.4	173/438	61.6 60.5	Total T	Matched 2:1	<sup>e</sup>	Bioavailable estradiol Age, storage time, subcohort, parity, and oophorectomy	2.1 (0.9–4.7) 1.9 (1.1–3.3)
Zeleniuch-Jacquotte, 2004 (101)	Excluded new cases diagnosed within 6 months of recruitment	<sup>e,h</sup>	297/563	60 60	Total T	Cohort	2.2 (1.3–3.6)	Estradiol, SHBG Age at menarche, family history of breast cancer, parity/age at first birth, history of total oophorectomy, history of breast cancer Estrone	1.9 (1.1–3.3) 2.4 (1.4–4.0)

T, Testosterone; N/A, not applicable because comparison was mean difference between cases and controls; RH, relative hazard; HT, hormone therapy; Not different, adjusting for these variables had no effect; BMI, body mass index.

<sup>a</sup> Effect ratios were presented as RR for Garland, Berrino, Dorgan, and Hankinson studies; as OR for Thomas and Zeleniuch-Jacquotte studies; and as RH for Cauley study.

<sup>b</sup> Effect ratios of total testosterone, except for Berrino and Cauley studies, in which effect ratios were for free testosterone.

<sup>c</sup> Mean age of all participants.

<sup>d</sup> No statistically significant difference of testosterone levels between cases and controls by mean comparison.

<sup>e</sup> Data not available.

<sup>f</sup> 95% CI was not available.

<sup>g</sup> Reanalysis after incident case was defined as at least 2-yr interval after blood taken resulted in RR = 1.3 (0.5–3.4).

<sup>h</sup> Median age at diagnosis was 66.1 yr.

reduced risk for breast cancer in young women in whom the number of GGC repeat lengths was greater than 17. In addition, they also suggested that AR repeat length (CAG or GGC) may be partly responsible for the increased risk for early-onset breast cancer in women who use oral contraceptives, although these findings need replication in other populations (116).

## VI. Aromatization and Breast Cancer Development

Estrogen biosynthesis in postmenopausal women is primarily the result of aromatization of circulating C19 steroids in extragonadal sites (1). The activity of the aromatase enzyme increases with age in fat tissue (117), and with increasing age there is greater fat tissue in the breast. Thus, hypothetically higher androgen levels in women provide increased substrate for estrogen biosynthesis within the breast. Despite this, it is also evident that estrogen itself can block its own bioformation in both human breast cancer cells and animal studies. Nakamura *et al.* (118) have shown that ovariectomy increases and estradiol treatment decreases aromatase activity in baboon mammary tissue. Subsequently, an inverse correlation between tumor aromatase activity and estrogen content has been reported in nude mice bearing xenografts of MCF-7 cells transfected with the aromatase gene (119). Moreover, an *in vitro* study in which MCF-7 cells were cultured long term in an estrogen-deprived medium and called by the acronym LTED cells found that long-term estrogen deprivation enhanced aromatase activity by 3- to 4-fold when compared with the wild-type MCF-7 cells (119). Reexposure of LTED cells to estrogen resulted in reducing aromatase activity to the levels of the wild-type MCF-7 cells (119). Consistent with these findings, intratumoral aromatase activity was higher in women with lower circulating estrogen (120, 121), and relative low activity was found in the patients taking hormone therapy (119, 122). These data suggest that after menopause, when circulating estrogen levels are low, an increase in aromatase levels in the breast may maintain tissue concentrations of estrogen. Thus, aromatase may control the local production of estrogen through an autocrine loop. During the process of transformation to malignancy, locally produced estrogen may stimulate the proliferation of tumor cells and vascular endothelial growth factor production. These effects are also likely to enhance tumor progression, development of angiogenesis, and, ultimately, metastasis of cancer. Therefore, it is more appropriate to use testosterone replacement only in women who have adequate estrogen replacement.

## VII. Should Androgens Be Included in Postmenopausal Hormone Therapy Regimens?

Whether there is a role for the use of androgens in the management of postmenopausal women remains controversial. Clinical studies of supraphysiological testosterone therapy have shown improvements in sexual parameters in postmenopausal women (123–125). More recent studies employing more physiological doses have shown benefits in several parameters of sexual function and in mood (126–128).

At present, a variety of testosterone-containing preparations are being used in clinical practice or in investigational research protocols for the treatment of sexual problems in women. Although the findings of this review indicate favorable effects of nonaromatizable androgens in the breast, in contrast to testosterone, there are few data from large well-designed randomized controlled trials to support the use of methyltestosterone in the management of sexual dysfunction in women (129). Also, data pertaining to the use of DHT in women are completely lacking. It is clear that whether or not the effects of testosterone on sexual function and mood in women are, in part, dependent on aromatization within the brain needs to be elucidated.

Combined oral estrogen-progestin postmenopausal therapy is associated with an increase in breast cancer risk (6, 7, 10). Whether this is an effect of oral estrogen or the inclusion of progestin is not known. However, the *in vitro* and *in vivo* data we have summarized indicate that in an estrogen-replete environment androgens oppose the unfavorable effects of estrogen in breast tissue. Consistent with this concept, tibolone, a synthetic steroid with estrogenic, progestogenic, and androgenic properties does not appear to have any adverse effects on breast tissue *in vitro* (130). In contrast, a large cohort study reported a RR of breast cancer of 1.45 (95% CI, 1.25–1.68) for users *vs.* nonusers of tibolone (131). However, these data should be viewed cautiously until reevaluated in a randomized controlled trial because of the inherent bias in the study population and the tendency for clinicians to prescribe tibolone rather than standard hormone therapy to women believed to be at increased breast cancer risk (132, 133). The Women's Health Initiative study reported that for the 12,304 women in the study who had never previously been treated with hormone therapy there was no increase in breast cancer risk for a mean duration of 5.2 yr of estrogen-progestin therapy (hazard ratio, 1.06; 95% CI, 0.81–1.28) (5–11). The significant increase in breast cancer risk reported in this study was related to the 4304 prior users (5–11). Thus, a large study would be required to demonstrate the risks *vs.* benefit of adding testosterone therapy to estrogen-progestin therapy for 5 yr; in addition, to explore any possibility that there may be a reduction in risk with testosterone, the study would need to be beyond 5 yr duration (133). An alternative approach would be to employ surrogate endpoints such as mammographic density and indices of breast cell proliferation and apoptosis. The latter requires breast needle biopsies in healthy women, a minor but invasive procedure. Although the findings from such research may be informative, there is still the danger of misleading results. An analogy is the substantial evidence that estrogen lowers lipids; yet in one large randomized controlled trial, estrogen-progestin therapy was associated with more cardiovascular events (5–11). Thus, with the available data pertaining to the effects of testosterone on the breast, the inclusion of testosterone in hormonal regimens should be limited to women symptomatic of androgen insufficiency despite adequate estrogen replacement. Testosterone therapy for women should involve regular measurements of circulating levels of free testosterone, and levels should be maintained below the upper limit of the normal physiological range for young women to avoid androgen excess (4).

### VIII. Conclusion

Breast cancer has complex etiologies; however, endogenous sex steroids clearly have a role in the progression of this disease. *In vitro* and *in vivo* studies indicate that both testosterone and DHT have a predominantly inhibitory influence on the mitogenic and cancer-promoting effects of estrogen in breast cells and promote apoptosis via the AR. There are, however, variations in these effects according to the type of breast cancer cell line studied, the androgen administered, and the dose used. These differences appear to be a consequence of differing levels of coactivator and corepressor proteins that influence AR actions in different cell types.

Unfortunately, most clinical studies have used total testosterone as a measure of androgen exposure, and these generally have shown that higher total testosterone levels are associated with increased breast cancer risk. However, these findings may reflect higher SHBG levels due to higher endogenous estrogen. There are few data pertaining to the relationship between free testosterone levels and breast cancer risk in humans using reliable assay methodology. Although studies in both premenopausal and postmenopausal women are inconclusive, there is no evidence that hyperandrogenism in women with PCOS is associated with increased breast cancer risk. Data for the use of exogenous testosterone and breast cancer risk are limited. The strongest supporting data for exogenous testosterone therapy come from primate studies. Based on such simulations, inclusion of testosterone in postmenopausal estrogen-progestin regimens has the potential to ameliorate the stimulating effects of combined estrogen-progestin on the breast. Research addressing this is warranted; however, the number of women that would be required for an adequately powered randomized controlled trial renders such a study unlikely. Unless more specific data become available, the use of testosterone should be limited to women symptomatic of androgen insufficiency despite adequate estrogen replacement.

### Acknowledgments

Address all correspondence and requests for reprints to: Professor Susan R. Davis, The Jean Hailes Foundation NHMRC Centre of Clinical Research Excellence, 173 Carinish Road, Clayton, Victoria 3168, Australia. E-mail: susan.davis@jeanhailes.org.au

### References

1. Simpson E, Rubin G, Clyne C, Robertson K, O'Donnell L, Jones M, Davis S 2000 The role of local estrogen biosynthesis in males and females. *Trends Endocrinol Metab* 11:184–188
2. Zumoff B, Strain GW, Miller LK, Rosner W 1995 Twenty-four hour mean plasma testosterone concentration declines with age in normal premenopausal women. *J Clin Endocrinol Metab* 80:1429–1430
3. Labrie F, Luu-The V, Labrie C, Belanger A, Simard J, Lin SX, Pelletier G 2003 Endocrine and intracrine sources of androgens in women: inhibition of breast cancer and other roles of androgens and their precursor dehydroepiandrosterone. *Endocr Rev* 24:152–182
4. Bachmann G, Bancroft J, Braunstein G, Burger H, Davis S, Dennerstein L, Goldstein I, Guay A, Leiblum S, Lobo R, Notelovitz M, Rosen R, Sarrel P, Sherwin B, Simon J, Simpson E, Shifren J, Spark R, Traish A 2002 Female androgen insufficiency: the Prince-
- ton consensus statement on definition, classification, and assessment. *Fertil Steril* 77:660–665
5. Persson I 2001 Estrogens in the causation of breast, endometrial and ovarian cancers—evidence and hypotheses from epidemiological findings. *J Steroid Biochem Mol Biol* 74:357–364
6. Ross R, Paganni Hill A, Wan P, Pike AC 2000 Effect of hormone replacement therapy on breast cancer risk; estrogen versus estrogen plus progestin. *J Natl Cancer Inst* 92:328–332
7. Schairer C, Lubin J, Troisi R, Sturgeon S, Brinton LA, Hoover R 2000 Menopausal estrogen and estrogen-progestin replacement therapy and breast cancer risk. *JAMA* 283:485–491
8. Beral V, Banks E, Reeves G 2002 Evidence from randomised trials on the long-term effects of hormone replacement therapy. *Lancet* 360:942–944
9. Magnusson C, Baron JA, Correia N, Bergstrom R, Adami HO, Persson I 1999 Breast-cancer risk following long-term oestrogen- and oestrogen-progestin-replacement therapy. *Int J Cancer* 81:339–344
10. Chlebowski RT, Hendrix SL, Langer RD, Stefanick ML, Gass M, Lane D, Rodabough RJ, Gilligan MA, Cyr MG, Thomson CA, Khandekar J, Petrovitch H, McTiernan A 2003 Influence of estrogen plus progestin on breast cancer and mammography in healthy postmenopausal women: the Women's Health Initiative randomized trial. *JAMA* 289:3243–3253
11. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, Jackson RD, Beresford SA, Howard BV, Johnson KC, Kotchen JM, Ockene J 2002 Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 288:321–333
12. Vaitukaitis JL, Dale SL, Melby JC 1969 Role of ACTH in the secretion of free DHA and its sulfate ester in man. *J Clin Endocrinol Metab* 29:1443–1447
13. Vermeulen A, Ando S 1978 Prolactin and adrenal androgen secretion. *Clin Endocrinol (Oxf)* 8:295–303
14. Haning Jr RV, Cabot M, Flood CA, Hackett R, Longcope C 1989 Metabolic clearance rate (MCR) of dehydroepiandrosterone sulfate (DS) its metabolism to dehydroepiandrosterone, androstenedione, testosterone and dihydrotestosterone, and the effects of increased plasma DS concentration on DS MCR in normal women. *J Clin Endocrinol Metab* 69:1047–1052
15. Davis SR 1999 Androgen replacement in women: a commentary. *J Clin Endocrinol Metab* 84:1886–1891
16. Burger HG 2002 Androgen production in women. *Fertil Steril* 77(Suppl 4):3–5
17. Simpson ER 2002 Aromatization of androgens in women: current concepts and findings. *Fertil Steril* 77(Suppl 4):6–10
18. Simpson E, Rubin G, Clyne C, Robertson K, O'Donnell L, Davis S, Jones M 1999 Local estrogen biosynthesis in males and females. *Endocr Relat Cancer* 6:131–137
19. Rannevik G, Jeppsson S, Johnell O, Bjerre B, Laurell-Borulf Y, Svanberg L 1995 A longitudinal study of the perimenopausal transition: altered profiles of steroid and pituitary hormones, SHBG and bone mineral density. *Maturitas* 21:103–113
20. Dunn JF, Nisula BC, Rodboard D 1981 Transport of steroid hormones. Binding of 21 endogenous steroids to both testosterone-binding globulin and cortico-steroid-binding globulin in human plasma. *J Clin Endocrinol Metab* 53:58–68
21. Davis S 2000 Androgens. In: Lobo R, Marcus R, Kelsey J, eds. *Menopause: biology and pathobiology*. San Diego, CA: Academic Press; 445–458
22. Mathur RS, Landgreve SC, Moody LO, Semmens JP, Williamson HO 1985 The effect of estrogen treatment on plasma concentrations of steroid hormones, gonadotropins, prolactin and sex hormone-binding globulin in post-menopausal women. *Maturitas* 7:129–133
23. Sinha-Hakim I, Arver S, Beall G, Shen R, Guerrero M, Sattler F, Shikuma C, Nelson JC, Landgren BM, Mazer NA, Bhasin S 1998 The use of a sensitive equilibrium dialysis method for the measurement of free testosterone levels in healthy, cycling women and in human immunodeficiency virus-infected women. *J Clin Endocrinol Metab* 83:1312–1318
24. Judd HL, Judd G, Lucas WE, Yen SSC 1974 Endocrine function of the postmenopausal ovary. Concentrations of androgens and es-

- trogens in ovarian and peripheral venous blood. *J Clin Endocrinol* 39:1020–1025
25. **Vierhapper H, Nowotny P, Waldhausl W** 1997 Determination of testosterone production rates in men and women using stable isotope dilution and mass spectrometry. *J Clin Endocrinol Metab* 82:1492–1496
  26. **Davis S, Schneider H, Donarti-Sarti C, Rees M, Van Lunsen H, Bouchard C, Derogatis L**, Androgen levels in normal and oophorectomized women. *Proc 10th International Congress on the Menopause*, Berlin, 2002, p 73 (Abstract F-12-01)
  27. **Mushayandebvu T, Castracane DV, Gimpel T, Adel T, Santoro N** 1996 Evidence for diminished midcycle ovarian androgen production in older reproductive aged women. *Fertil Steril* 65:721–723
  28. **Longcope C, Franz C, Morello C, Baker K, Johnston Jr CC** 1986 Steroid and gonadotropin levels in women during the peri-menopausal years. *Maturitas* 8:189–196
  29. **Judd HL, Lucas WE, Yen SSC** 1994 Effect of oophorectomy on circulating testosterone and androstenedione levels in patients with endometrial cancer. *Am J Obstet Gynecol* 118:793–798
  30. **Couzinet B, Meduri G, Lecce M, Young J, Brailly S, Loosfelt H, Milgrom E, Schaison G** 2001 The postmenopausal ovary is not a major androgen-producing gland. *J Clin Endocrinol Metab* 86:5060–5065
  31. **Zhou J, Ng S, Adesanya-Famuyi O, Anderson K, Bondy CA** 2000 Testosterone inhibits estrogen-induced mammary epithelial proliferation and suppresses estrogen receptor expression. *FASEB J* 14:1725–1730
  32. **Dimitrakakis C, Zhou J, Bondy CA** 2002 Androgens and mammary growth and neoplasia. *Fertil Steril* 77:S26–S33
  33. **Kerr JF, Wyllie AH, Currie AR** 1972 Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br J Cancer* 26:239–257
  34. **Feuerhake F, Sigg W, Hofter EA, Unterberger P, Welsch U** 2003 Cell proliferation, apoptosis, and expression of Bcl-2 and Bax in non-lactating human breast epithelium in relation to the menstrual cycle and reproductive history. *Breast Cancer Res Treat* 77:37–48
  35. **Ferguson DJ, Anderson TJ** 1981 Morphological evaluation of cell turnover in relation to the menstrual cycle in the “resting” human breast. *Br J Cancer* 44:177–181
  36. **Daniel CW, Silberstein GB, Strickland P** 1987 Direct action of 17 $\beta$ -estradiol on mouse mammary ducts analyzed by sustained release implants and steroid autoradiography. *Cancer Res* 47:6052–6057
  37. **Haslam SZ** 1988 Progesterone effects on deoxyribonucleic acid synthesis in normal mouse mammary glands. *Endocrinology* 122:464–470
  38. **Silberstein GB, Van Horn K, Shyamala G, Daniel CW** 1994 Essential role of endogenous estrogen in directly stimulating mammary growth demonstrated by implants containing pure antiestrogens. *Endocrinology* 134:84–90
  39. **Massafra C, De Felice C, Agnusdei DP, Gioia D, Bagnoli F** 1999 Androgens and osteocalcin during the menstrual cycle. *J Clin Endocrinol Metab* 84:971–974
  40. **Anderson E** 2001 Ovarian steroids and control of proliferation in the normal human breast. *Breast* 10:273–278
  41. **Malet C, Gompel A, Yaneva H, Cren H, Fidji N, Mowszowicz J, Kuttann F, Mauvais-Jarvis P** 1991 Estradiol and progesterone receptors in cultured normal human breast epithelial cells and fibroblasts: immunocytochemical studies. *J Clin Endocrinol Metab* 73:8–17
  42. **Clarke RB, Laidlaw IJ, Jones LJ, Howell A, Anderson E** 1993 Effect of tamoxifen on Ki67 labelling index in human breast tumours and its relationship to oestrogen and progesterone receptor status. *Br J Cancer* 67:606–611
  43. **Clarke RB, Howell A, Anderson E** 1997 Estrogen sensitivity of normal human breast tissue in vivo and implanted into athymic nude mice: analysis of the relationship between estrogen-induced proliferation and progesterone receptor expression. *Breast Cancer Res Treat* 45:121–133
  44. **Laidlaw IJ, Clarke RB, Howell A, Owen AW, Potten CS, Anderson E** 1995 The proliferation of normal human breast tissue implanted into athymic nude mice is stimulated by estrogen but not progesterone. *Endocrinology* 136:164–171
  45. **Kyprianou N, English HF, Davidson NE, Isaacs JT** 1991 Programmed cell death during regression of the MCF-7 human breast cancer following estrogen ablation. *Cancer Res* 51:162–166
  46. **Perry RR, Kang Y, Greaves B** 1995 Effects of tamoxifen on growth and apoptosis of estrogen-dependent and -independent human breast cancer cells. *Ann Surg Oncol* 2:238–245
  47. **Graham JD, Clarke CL** 1997 Physiological action of progesterone in target tissues. *Endocr Rev* 18:502–519
  48. **Ando S, De Amicis F, Rago V, Carpino A, Maggiolino M, Panno M, Lanzino M** 2002 Breast cancer: from estrogen to androgen receptor. *Mol Cell Endocrinol* 193:121–125
  49. **Wang DY, Allen DS, De Stavola BL, Fentiman IS, Brussen J, Bulbrook RD, Thomas BS, Hayward JL, Reed MJ** 2000 Urinary androgens and breast cancer risk: results from a long-term prospective study based in Guernsey. *Br J Cancer* 82:1577–1584
  50. **Birrell SN, Roder DM, Horsfall DJ, Bentel JM, Tilley WD** 1995 Medroxyprogesterone acetate therapy in advanced breast cancer: the predictive value of androgen receptor expression. *J Clin Oncol* 13:1572–1577
  51. **Bocuzzi G, Brignardello E, Di Monaco M, Gatto V, Leonardi L, Pizzini A, Gallo M** 1994 5-En-androstene-3 $\beta$ ,17 $\beta$ -diol inhibits the growth of MCF-7 breast cancer cells when oestrogen receptors are blocked by oestradiol. *Br J Cancer* 70:1035–1039
  52. **Birrell SN, Bentel JM, Hickey TE, Ricciardelli C, Weger MA, Horsfall DJ, Tilley WD** 1995 Androgens induce divergent proliferative responses in human breast cancer cell lines. *J Steroid Biochem Mol Biol* 52:459–467
  53. **Birrell SN, Hall RE, Tilley WD** 1998 Role of the androgen receptor in human breast cancer. *J Mammary Gland Biol Neoplasia* 3:95–103
  54. **Ortmann J, Prifti S, Bohlmann MK, Rehberger-Schneider S, Strowitzki T, Rabe T** 2002 Testosterone and 5 $\alpha$ -dihydrotestosterone inhibit in vitro growth of human breast cancer cell lines. *Gynecol Endocrinol* 16:113–120
  55. **Magklara A, Brown TJ, Diamandis EP** 2002 Characterization of androgen receptor and nuclear receptor co-regulator expression in human breast cancer cell lines exhibiting differential regulation of kallikreins 2 and 3. *Int J Cancer* 100:507–514
  56. **Yu H, Giai M, Diamandis EP, Katsaros D, Sutherland DJ, Levesque MA, Roagna R, Ponzone R, Sismondi P** 1995 Prostate-specific antigen is a new favorable prognostic indicator for women with breast cancer. *Cancer Res* 55:2104–2110
  57. **Magklara A, Grass L, Diamandis EP** 2000 Differential steroid hormone regulation of human glandular kallikrein (hK2) and prostate-specific antigen (PSA) in breast cancer cell lines. *Breast Cancer Res Treat* 59:263–270
  58. **Kandouz M, Lombet A, Perrot JY, Jacob D, Carvajal S, Kazem A, Rostene W, Therwath A, Gompel A** 1999 Proapoptotic effects of antiestrogens, progestins and androgen in breast cancer cells. *J Steroid Biochem Mol Biol* 69:463–471
  59. **Herrmann J, Bruckheimer E, McDonnell T** 1996 Cell death signal transduction and bcl-2 function. *Biochem Soc Trans* 24:1059–1065
  60. **Xie B, Tsao SW, Wong YC** 1999 Sex hormone-induced mammary carcinogenesis in female noble rats: the role of androgens. *Carcinogenesis* 20:1597–1606
  61. **Lapointe J, Fournier A, Richard V, Labrie C** 1999 Androgens down-regulate bcl-2 protooncogene expression in ZR-75-1 human breast cancer cells. *Endocrinology* 140:416–421
  62. **Xie B, Tsao SW, Wong YC** 2000 Sex hormone-induced mammary carcinogenesis in the female Noble rats: expression of bcl-2 and bax in hormonal mammary carcinogenesis. *Breast Cancer Res Treat* 61:45–57
  63. **Li S, Yan X, Belanger A, Labrie F** 1994 Prevention by dehydroepiandrosterone of the development of mammary carcinoma induced by 7,12-dimethylbenz(a)anthracene (DMBA) in the rat. *Breast Cancer Res Treat* 29:203–217
  64. **Luo S, Labrie C, Belanger A, Labrie F** 1997 Effect of dehydroepiandrosterone on bone mass, serum lipids, and dimethylbenz(a)anthracene-induced mammary carcinoma in the rat. *Endocrinology* 138:3387–3394
  65. **Couillard S, Labrie C, Belanger A, Candas B, Pouliot F, Labrie F** 1998 Effect of dehydroepiandrosterone and the antiestrogen EM-800 on growth of human ZR-75-1 breast cancer xenografts. *J Natl Cancer Inst* 90:772–778

66. Lubet RA, Gordon GB, Prough RA, Lei XD, You M, Wang Y, Grubbs CJ, Steele VE, Kelloff GJ, Thomas CF, Moon RD 1998 Modulation of methylnitrosourea-induced breast cancer in Sprague Dawley rats by dehydroepiandrosterone: dose-dependent inhibition, effects of limited exposure, effects on peroxisomal enzymes, and lack of effects on levels of Ha-Ras mutations. *Cancer Res* 58:921–926
67. Dauvois S, Li SM, Martel C, Labrie F 1989 Inhibitory effect of androgens on DMBA-induced mammary carcinoma in the rat. *Breast Cancer Res Treat* 14:299–306
68. Dauvois S, Labrie F 1989 Androstenedione and androst-5-ene-3 $\beta$ ,17 $\beta$ -diol stimulate DMBA-induced rat mammary tumors—role of aromatase. *Breast Cancer Res Treat* 13:61–69
69. Dauvois S, Geng CS, Levesque C, Merand Y, Labrie F 1991 Additive inhibitory effects of an androgen and the antiestrogen EM-170 on estradiol-stimulated growth of human ZR-75-1 breast tumors in athymic mice. *Cancer Res* 51:3131–3135
70. Dimitrakakis C, Zhou J, Wang J, Belanger A, Labrie F, Cheng C, Powell D, Bondy C 2003 A physiologic role for testosterone in limiting estrogenic stimulation of the breast. *Menopause* 10:292–298
71. Jayo MJ, Register TC, Hughes CL, Blas-Machado U, Sulistiawati E, Borgerink H, Johnson CS 2000 Effects of an oral contraceptive combination with or without androgen on mammary tissues: a study in rats. *J Soc Gynecol Invest* 7:257–265
72. Sourla A, Martel C, Labrie C, Labrie F 1998 Almost exclusive androgenic action of dehydroepiandrosterone in the rat mammary gland. *Endocrinology* 139:753–764
73. Baird DT, Horton R, Longcope C, Tait JF 1969 Steroid dynamics under steady-state conditions. *Recent Prog Horm Res* 25:611–664
74. Vermeulen A, Rubens R, Verdonck L 1972 Testosterone secretion and metabolism in male senescence. *J Clin Endocrinol Metab* 34:730–735
75. Batrinis ML, Panitsa-Faflija C, Koutsoumanis C, Vourlioti T, Koutsilieris M 1999 Surgical stress induces a marked and sustained increase of adrenal androgen secretion in postmenopausal women. *In Vivo* 13:147–150
76. Calogero AE, Burrello N, Negri-Cesi P, Papale L, Palumbo MA, Cianci A, Sanfilippo S, D'Agata R 1996 Effects of corticotropin-releasing hormone on ovarian estrogen production *in vitro*. *Endocrinology* 137:4161–4166
77. Calogero AE, Barreca A, Burrello N, Palermo I, Giordano G, D'Agata R, Vicari E 2002 Corticotrophin-releasing hormone inhibits insulin-like growth factor-I release from primary cultures of rat granulosa cells. *J Endocrinol* 174:493–498
78. Cruess D, Antoni M, Kumar M, McGregor B, Alferi S, Boyers A, Carver C, Kilbourn K 2001 Effects of stress management on testosterone levels in women with early-stage breast cancer. *Int J Behav Med* 8:194–207
79. Labrie F, Luu-The V, Lin SX, Simard J, Labrie C, El Alfy M, Pelletier G, Belanger A 2000 Intracrinology: role of the family of 17  $\beta$ -hydroxysteroid dehydrogenases in human physiology and disease. *J Mol Endocrinol* 25:1–16
80. Labrie F 2003 Extragonadal synthesis of sex steroids: intracrinology. *Ann Endocrinol (Paris)* 64:95–107
81. Thomas HV, Key TJ, Allen DS, Moore JW, Dowsett M, Fentiman IS, Wang DY 1997 A prospective study of endogenous serum hormone concentrations and breast cancer risk in post-menopausal women on the island of Guernsey. *Br J Cancer* 76:401–405
82. Garland CF, Friedlander NJ, Barrett-Connor E, Khaw KT 1992 Sex hormones and postmenopausal breast cancer: a prospective study in an adult community. *Am J Epidemiol* 135:1220–1230
83. Klee GG, Hesser D 2000 Techniques to measure testosterone in the elderly. *Mayo Clin Proc* 75:S19–S25
84. Secreto G, Toniolo P, Pisani P, Recchione C, Cavalleri A, Fariselli G, Totis A, Di Pietro S, Berrino F 1989 Androgens and breast cancer in premenopausal women. *Cancer Res* 49:471–476
85. Yu H, Shu XO, Shi R, Dai Q, Jin F, Gao YT, Li BD, Zheng W 2003 Plasma sex steroid hormones and breast cancer risk in Chinese women. *Int J Cancer* 105:92–97
86. Wysowski DK, Comstock GW, Helsing KJ, Lau HL 1987 Sex hormone levels in serum in relation to the development of breast cancer. *Am J Epidemiol* 125:791–799
87. Thomas HV, Key TJ, Allen DS, Moore JW, Dowsett M, Fentiman IS, Wang DY 1997 A prospective study of endogenous serum hormone concentrations and breast cancer risk in premenopausal women on the island of Guernsey. *Br J Cancer* 75:1075–1079
88. Fox R, Corrigan E, Thomas PG, Hull MG 1991 Oestrogen and androgen states in oligo-amenorrhoeic women with polycystic ovaries. *Br J Obstet Gynaecol* 98:294–299
89. Coulam CB, Annegers JF, Kranz JS 1983 Chronic anovulation syndrome and associated neoplasia. *Obstet Gynecol* 61:403–407
90. Anderson KE, Sellers TA, Chen PL, Rich SS, Hong CP, Folsom AR 1997 Association of Stein-Leventhal syndrome with the incidence of postmenopausal breast carcinoma in a large prospective study of women in Iowa. *Cancer* 79:494–499
91. Gammon MD, Thompson WD 1991 Polycystic ovaries and the risk of breast cancer. *Am J Epidemiol* 134:818–824
92. Lipworth L, Adami HO, Trichopoulos D, Carlstrom K, Mantzoros C 1996 Serum steroid hormone levels, sex hormone-binding globulin, and body mass index in the etiology of postmenopausal breast cancer. *Epidemiology* 7:96–100
93. Secreto G, Toniolo P, Berrino F, Recchione C, Cavalleri A, Pisani P, Totis A, Fariselli G, Di Pietro S 1991 Serum and urinary androgens and risk of breast cancer in postmenopausal women. *Cancer Res* 51:2572–2576
94. Grattarola R, Secreto G, Recchione C, Castellini W 1974 Androgens in breast cancer. II. Endometrial adenocarcinoma and breast cancer in married postmenopausal women. *Am J Obstet Gynecol* 118:173–178
95. Berrino F, Muti P, Michelli A, Bolelli G, Krogh V, Sciajno R, Pisani P, Panico S, Secreto G 1996 Serum sex hormone levels after menopause and subsequent breast cancer. *J Natl Cancer Inst* 88:291–296
96. Dorgan JF, Longcope C, Stephenson Jr HE, Falk RT, Miller R, Franz C, Kahle L, Campbell WS, Tangrea JA, Schatzkin A 1996 Relation of prediagnostic serum estrogen and androgen levels to breast cancer risk. *Cancer Epidemiol Biomarkers Prev* 5:533–539
97. Zeleniuch-Jacquotte A, Bruning PF, Bonfrer J, Koenig K, Shore R, Kim M, Pasternack B, Toniolo P 1997 Relation of serum levels of testosterone and dehydroepiandrosterone sulfate to risk of breast cancer in postmenopausal women. *Am J Epidemiol* 145:1030–1038
98. Hankinson SE, Willett WC, Manson JE, Colditz GA, Hunter DJ, Spiegelman D, Barbieri R, Speizer FE 1998 Plasma sex steroid hormone levels and risk of breast cancer in postmenopausal women. *J Natl Cancer Inst* 90:1292–1299
99. Cauley JA, Lucas FL, Kuller LH, Stone K, Browner W, Cummings SR 1999 Elevated serum estradiol and testosterone concentrations are associated with a high risk for breast cancer. Study of Osteoporotic Fractures Research Group. *Ann Intern Med* 130:270–277
100. Manjer J, Johansson R, Berglund G, Janzon L, Kaaks R, Agren A, Lenner P 2003 Postmenopausal breast cancer risk in relation to sex steroid hormones, prolactin and SHBG (Sweden). *Cancer Causes Control* 14:599–607
101. Zeleniuch-Jacquotte A, Shore RE, Koenig KL, Akhmedkhanov A, Afanasyeva Y, Kato I, Kim MY, Rinaldi S, Kaaks R, Toniolo P 2004 Postmenopausal levels of oestrogen, androgen, and SHBG and breast cancer: long-term results of a prospective study. *Br J Cancer* 90:153–159
102. Hankinson SE, Manson JE, Spiegelman D, Willett WC, Longcope C, Speizer FE 1995 Reproducibility of plasma hormone levels in postmenopausal women over a 2–3-year period. *Cancer Epidemiol Biomarkers Prev* 4:649–654
103. Endogenous Hormone and Breast Cancer Collaborative Group 2002 Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 94:606–616
104. Barrett-Connor E, Friedlander NJ, Khaw KT 1990 Dehydroepiandrosterone sulfate and breast cancer risk. *Cancer Res* 50:6571–6574
105. Lonning PE 1996 Aromatase inhibition for breast cancer treatment. *Acta Oncol* 35(Suppl 5):38–43
106. Moehrer B, Hextall A, Jackson S 2003 Oestrogens for urinary incontinence in women. *Cochrane Database Syst Rev* CD001405
107. Dimitrakakis C, Jones R, Liu A, Bondy C 2003 Breast cancer incidence

- in Australian women using testosterone in addition to estrogen replacement. Program of the 85th Annual Meeting of The Endocrine Society, Philadelphia, 2003, p 574 (Abstract P3-424)
108. **Brinton LA, Hoover R, Fraumeni Jr JF** 1986 Menopausal oestrogens, and breast cancer risk: an expanded study case-control study. *Br J Cancer* 54:825–832
  109. **Ewertz M** 1988 Influence of non-contraceptive exogenous and endogenous sex hormones on breast cancer risk in Denmark. *Int J Cancer* 42:832–838
  110. **Recchione C, Venturelli E, Manzari A, Cavalteri A, Martinetti A, Secreto G** 1995 Testosterone, dihydrotestosterone and estradiol levels in postmenopausal breast cancer tissues. *J Steroid Biochem Mol Biol* 52:541–546
  111. **Soreide JA, Lea OA, Varhaug JE, Skarstein A, Kvinnsland S** 1992 Androgen receptors in operable breast cancer: relation to other steroid hormone receptors, correlations to prognostic factors and predictive value for effect of adjuvant tamoxifen treatment. *Eur J Surg Oncol* 18:112–118
  112. **Selim AG, El Ayat G, Wells CA** 2002 Androgen receptor expression in ductal carcinoma in situ of the breast: relation to oestrogen and progesterone receptors. *J Clin Pathol* 55:14–16
  113. **Chamberlain NL, Driver ED, Miesfeld RL** 1994 The length and location of CAG trinucleotide repeats in the androgen receptor N-terminal domain affect transactivation function. *Nucleic Acids Res* 22:3181–3186
  114. **Rebeck TR, Kantoff PW, Krithivas K, Neuhausen S, Blackwood MA, Godwin AK, Daly MB, Narod SA, Garber JE, Lynch HT, Weber BL, Brown M** 1999 Modification of BRCA1-associated breast cancer risk by the polymorphic androgen-receptor CAG repeat. *Am J Hum Genet* 64:1371–1377
  115. **Kadouri L, Easton DF, Edwards S, Hubert A, Kote-Jarai Z, Glaser B, Durocher F, Abeliovich D, Peretz T, Eeles RA** 2001 CAG and GGC repeat polymorphisms in the androgen receptor gene and breast cancer susceptibility in BRCA1/2 carriers and non-carriers. *Br J Cancer* 85:36–40
  116. **Suter NM, Malone KE, Daling JR, Doody DR, Ostrander EA** 2003 Androgen receptor (CAG)(n) and (GGC)(n) polymorphisms and breast cancer risk in a population-based case-control study of young women. *Cancer Epidemiol Biomarkers Prev* 12:127–135
  117. **Bulun SE, Simpson ER** 1994 Competitive R. T-PCR analysis indicates levels of aromatase cytochrome P450 transcripts in adipose tissue of buttocks, thighs and abdomen of women increase with advancing age. *J Clin Endocrinol Metab* 78:428–432
  118. **Nakamura J, Lu Q, Aberdeen G, Albrecht E, Brodie A** 1999 The effect of estrogen on aromatase and vascular endothelial growth factor messenger ribonucleic acid in the normal nonhuman primate mammary gland. *J Clin Endocrinol Metab* 84:1432–1437
  119. **Yue W, Berstein LM, Wang JP, Clark GM, Hamilton CJ, Demers LM, Santen RJ** 2001 The potential role of estrogen in aromatase regulation in the breast. *J Steroid Biochem Mol Biol* 79:157–164
  120. **Berstein LM, Larionov AA, Kyshtoobaeva AS, Pozharisski KM, Semiglazov VF, Ivanova OA** 1996 Aromatase in breast cancer tissue—localization and relationship with reproductive status of patients. *J Cancer Res Clin Oncol* 122:495–498
  121. **Bolufer P, Ricart E, Lluch A, Vazquez C, Rodriguez A, Ruiz A, Llopis F, Garcia-Conde J, Romero R** 1992 Aromatase activity and estradiol in human breast cancer: its relationship to estradiol and epidermal growth factor receptors and to tumor-node-metastasis staging. *J Clin Oncol* 10:438–446
  122. **Pasqualini JR, Chetrite G, Blacker C, Feinstein MC, Delalonde L, Talbi M, Maloche C** 1996 Concentrations of estrone, estradiol, and estrone sulfate and evaluation of sulfatase and aromatase activities in pre- and postmenopausal breast cancer patients. *J Clin Endocrinol Metab* 81:1460–1464
  123. **Burger H, Hailes J, Nelson J, Menelaus M** 1987 Effect of combined implants of oestradiol and testosterone on libido in postmenopausal women. *Br Med J (Clin Res Ed)* 294:936–937
  124. **Sherwin BB, Gelfand MM, Brender W** 1985 Androgen enhances sexual motivation in females: a prospective, crossover study of sex steroid administration in surgical menopause. *Psychosom Med* 47:339–351
  125. **Studd JWW, Colins WP, Chakravarti S** 1977 Estradiol and testosterone implants in the treatment of psychosexual problems in postmenopausal women. *Br J Obstet Gynaecol* 84:314–315
  126. **Davis SR, McCloud P, Strauss BJ, Burger H** 1995 Testosterone enhances estradiol's effects on postmenopausal bone density and sexuality. *Maturitas* 21:227–236
  127. **Shifren JL, Braunstein G, Simon J, Casson P, Buster JE, Red Burki RE, Ginsburg ES, Rosen RC, Leiblum SR, Caramelli KE** 2000 Transdermal testosterone treatment in women with impaired sexual function after oophorectomy. *N Engl J Med* 343:682–688
  128. **Goldstat R, Briganti E, Tran J, Wolfe R, Davis SR** 2003 Transdermal testosterone therapy improves well-being, mood, and sexual function in premenopausal women. *Menopause* 10:390–398
  129. **Lobo RA, Rosen RC, Yang HM, Block B, Van Der Hoop RG** 2003 Comparative effects of oral esterified estrogens with and without methyltestosterone on endocrine profiles and dimensions of sexual function in postmenopausal women with hypoactive sexual desire. *Fertil Steril* 79:1341–1352
  130. **Kloosterboer H** 2001 Tibolone: a steroid with tissue-specific mode of action. *J Steroid Biochem Mol Biol* 76:231–238
  131. **Beral V** 2003 Breast cancer and hormone-replacement therapy in the Million Women Study. *Lancet* 362:419–427
  132. **Million Women Study Collaborators** 2002 Patterns of use of hormone replacement therapy in one million women in Britain 1996–2000. *Br J Obstet Gynaecol* 109:1319–1330
  133. **Collaborative Group on Hormonal Factors in Breast Cancer** 1997 Breast cancer and hormone replacement therapy: collaborative re-analysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. *Lancet* 350:1047–1059

*Endocrine Reviews* is published bimonthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.